PINADZHYAN, V.V.; INDZHIKYAN, Ya A

Deformation of plastic steel under the combined effect of stretching and torsion. Izv.AN Arm. SSR. Ser. tekh. nauk. 12 no.1:53-56 159. (MIRA 12:4)

1. Institut stroymaterialov i soorusheniy Ministerstva stroitel - stva Arm. SSR.

(Steel--Testing)

(Deformation (Mechanics))

APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-0

JNFRBATEV, M.S.

Errors of the method of difference for second-order elliptic equations. Vest. AN Kazakh. SSR 19 no.11.93-96 Nº53.. (HJRA 17:5)

INERBAYEV, M.S.

Errors of difference solutions to the second and third boundary value problems for elliptic equations. Metod. vych. no.2:50-59 '63. (MIRA 18:11)

INES, Z.

New safety measures for high-tension electric cable network in mines. p. 115.

PRZEGIAD GORNICZY. Stowarzyszenie Naukowo-Techniczne Inzynierow i Technikow Gornictwa. Katowice, Poland, Vol. 15, No. 3, March, 1959.

Monthly List of East European Accessions (EEAI), IC, Vol. 8, No. 9, September, 1959. Uncl.

5/271/63/000/003/012 A060/1126

Avraamov, I.S., Ineshin, A.P.

TITLE:

Engineering logic and the automation of production

PERIODICAL: Referativnyy zhurnal, Avtomatika, telemekhunika i vychislitel naya tekhnika, no. 3, 1963, 55, abstract 3A312 Woh. sap. Tomskiy un-t, 1962, no. 41, 156 - 170)

The authors describe a digital servosystem designed for controlling TEXT: a large class of rachanisms connected with the displacement and precise stopping at various points. To such mechanisms belongs: factory cranes, pressure units of rolling mills, mine elevators, ingot cars, etc. The system contains a memory unit for the coordinates of the exact technical stopping point, a memony unit of the current position of mechanisms, a feedback transducer and demputer unit. With the aid of the methods of the algebra of logic a reliable computer metwork is worked out. The reliability of its operation is attained through the application of a reflecting code, the introduction of DC feedbacks and of stabilizing networks which protect the flip-flops from pulse noise. There are 9 figures and

Card 1/2

		1.	The second of the		
Engineering logic and	the automation of pro	duction	3/271/63/000, 4060/4126	(003/012/049	
4 references.					
[Abstractor's note:	Complete translation]		A. S.		
Card 2/2					
	G .				

L 05410-67 EMT(d)/EMP(V)/EMP(h)/EMT(1)

ACC NR: AT6022758 SOURCE CODE: UR/2563/65/000/259/0107/0114

AUTHOR: Drannikov, V. G.; Yesin, A. I.; Ineshin, A. P.; Sevast'yanov, V. A.

33

ORG: None

TITLE: Analysis of the dynamics of a self-saturating magamp drive with intermediate semiconductor amplifiers

SOURCE: Leningrad. Politekhnicheskiy institut. Trudy, no. 259, 1965. Perekhodnyye protsessy v avtomatizirovannom elektroprivode (Transient processes in automated electric drive), 107-114

TOPIC TAGS: magnetic amplifier, machine tool, industrial automation

ABSTRACT: The authors consider the use of intermediate semiconductor amplifiers as a means for reducing the time constant in self-saturating magnetic-amplifier circuits used in combination with electric motors for driving the <u>feed</u> screws of machine tools. An analysis of transition processes in this type of system shows that linear operation of the intermediate semiconductor amplifier in self-saturating magamp-motor drive combinations has no noticeable effect on the time constant of the drive. The interference voltage acting through the correction circuit in an actual drive puts the intermediate amplifier into conditions of artificial switching with a frequency of 300 cps which increases the time of the transition process by a factor of more than 1.5. Class D

Card 1/2

APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3'


ACC NR: AT6022758

intermediate semiconductor rectifiers with pdm may be used satisfactorily for widerange control in self-saturating magamp drives. The small losses in the output tranrange control in self-saturating magamp drives. The small losses in the output transistor of the amplifier in both the open and closed states result in considerable
power delivery at high efficiency to the control circuits of the magnetic amplifier.
power delivery at high efficiency to the control circuits of the scatter in its paraThe operation of this transistor is nearly independent of the scatter in its parameters and variations in ambinet temperature. The frequency of the intermediate amplifier must be selected with regard to the particular features of the specific magplifier must be selected with regard to the particular features of the specific magplifier must be selected with regard to the particular features of the specific magplifier with high-power class D intermediate semiconductor amplifiers provides highjunction with high-power class D intermediate semiconductor amplifiers provides highjunction with high-power class D intermediate semiconductor amplifiers provides highjunction with high-power class D intermediate semiconductor amplifiers provides highjunction vith high-power class D intermediate semiconductor amplifiers provides highjunction vith high-power class D intermediate semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor amplifiers provides highjunction vith high-power class D intermediate, semiconductor

Card 2/2 - 14

APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3



NESHIND, 1.11.

Category: USSR/Analytical Chemistry - General Questions.

G-1

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 30945

Author : Ginzburg V. L., Alekseyenko Ye. F., Belokrinitakaya Ye. Ye.,

Vitushkina I. N., Ineshina F. M.

Inst: : not given

Title : Accuracy of Photographic Methods of Spectral Analysis

Orig Pub: Zavod. laboratoriya, 1956, 22, No 11, 1331-1333

Abstract: A comparison was made of the accuracy of analyses of fused nickel, copper regulus, fused cobalt and cathodic nickel, according to calibration graphs in \(\triangle \) S, lg C coordinates, and in accordance with the solid graph method. Determinations were made of Cu, Fe, Au, Pt, Pd, Ni, Si, Mn, Pb, Sb, Bi, Sn, Co, at concentrations from several thousandth to decimal fractions of one percent, with spectrum excitation in arc discharge of direct and alternating current, and photographic recording on plates of type I, II and III. In most instances no substantial differences were found in the magnitude of errors with different calibration graphs.


Card : 1/1

-18-

APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-00513R00061861001

"APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3



APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3

THE PERSON OF TH

PHASE I BOOK EXPLOITATION

30V/6260

Gurvich, Lev Veniaminovich, Georgiy Akopovich Khachkuruzov, Vadim Andreyevich Medvedev, Inessa Veniaminovna Veyts, Georgiy Andreyevich Bergman, Vladimir Stepanovich Yungman, Nina Petrovna Rtishcheva, Lidiya Fedorovna Kuratova, Georgiy Nikolayevich Yurkov, Amaliya Abramovna Kane, Boris Fedorovich Yucki, Boris Isidorovich Brounshteyn, Viktor Feodoseyevich Baybuz, Valeriy Aleksandrovich Kvlividze, Yevgeniy Aleksandrovich Prozorovskiy, and Boris Aleksandrovich Vorob'yev.

Termodinamicheskiye svoystva individual'nykh veshchestv; spravochnik v dvukh tomakh. tom 1: Vychisleniye termodinamicheskikh svoystv; tom 2: Tablitsy termodinamicheskikh svoystv (Thermodynamic Properties of Individual Substances; Reference Book in Two Volumes. v. 1: Calculation of Thermodynamic Properties; v. 2: Tables of Thermodynamic Properties). 2d ed., rev. and enl. Moscow, Izd-vo AN SSSR, 1962. 1161 and 916 p. 4000 copies printed.

Sponsoring Agencies: Akademiya nauk SSSR. Institut goryuchikh iskopayemykh; and Gosudarstvennyy komitet Soveta Ministrov SSSR Card 1/9/2

po khimii. Institut prikladnoy khimii.

Resp. Ed.: V. P. Glushko, Academician, L. V. Gurvich, G. A. Khach-kuruzov, I. V. Veyts, and V. A. Medvedev; Ed. of Publishing House: K. P. Gurov; Tech. Ed.: V. G. Laut.

PURPOSE: This reference book may be used in scientific-research and experimental-design work in institutes, design offices, and schools of higher education, as well as for training specialists in chemical thermodynamics and thermal physics.

COVERAGE: Volume 1 of this work deals with methods for calculating thermodynamic properties and with the selection of constants required for the calculations. Volume 2 contains tables of thermodynamic properties (reduced thermodynamic potential, entropy, enthalpy, and the logarithm of the dissociation or ionization constants of equilibrium) compiled, where data were lacking, on the basis of published and unpublished material from a number of Soviet research institutes. Thermodynamic properties for the ideal gas

Card 2/8/3

Thermodynamic Properties (Cont.)

SOV/6260

state are presented in table form for 335 gases, 44 liquids, and 45 solids compounded from 33 chemical elements and their isotopes, viz.: H, D, T, He, Li, Be, B, C, N, O, F, Ne, Na, Mg, Al, Si, P, S, Cl, Ar, K, Ca, Br, Kr, Re, Sr, Zr, I, Xe, Cs, Ba, Hg, and Pb. Thermodynamic properties are given for the following 22 gases in the range from room temperature to 20,000 K: H,H', H'', O, O', Na, O', O', O', O', Na, No, NO', C, C', CO, CO', and e'; for the 14 least stable gases up to 4000 K; and for the remaining 299 gases up to 6000 K. Virial coefficients for 34 gases are also given up to 6000 K.

TABLE OF CONTENTS (Volume 1) [Abridged]:

Foreword

Introduction

11

19

PART I. METHODS OF CALCULATING THE THERMODYNAMIC PROPERTIES OF INDIVIDUAL SUBSTANCES

Card 3/8/3

INEV, V.

INEV, V. Improving the technological work in shunting is an important condition for reducing the stopover of the railroad cars. p. 8. Vol. 8, no. 6, 1956. TRANSPORTNO DELO. Sofiia, Bulgaria

SOURCE: East European Accessions List (EEAL) Vol 6, No. 4-April 1957

INFANT'YEV, A.A., gornyy inzh., MITROFANCV, A.I., gornyy inzh.

Experience in desp dreinnge at the Yakovleve mine in the Kursk Megnetic Anomaly. Gor. zhur. no.il.ile.22 N '63.

1. Yakowlevskiy rudnik Kurskoy magnitnoy anomalii.

(MIRA 17:6)

18(5),14(5)

AUTHORS:

SOV/127-59-2-3/21 Gusev, A.M., Red'ko, L.A., and Infant'yev, A.N.

Mining Engineers

TITLE:

Preliminary Considerations Concerning the Methods of Opening, and Ways of Mining in the Yakovleveleye Deposit Area (Proyektnyye soobrazheniya o metodakh vskrytiya i sposobakh razrabotki Yakovlevskogo mesto-

arendinario in series de la companio del companio de la companio de la companio del companio de la companio del companio de la companio de la companio de la companio de la companio del companio de la companio del companio del companio del companio de la companio de la companio del companio d

PERIODICAL:

Gornyy zhurnal, 1959, Nr 2, pp 10-15 (USSR)

ABSTRACT:

The authors first give a concise description of the Yakovlevskoye and Pokrovskoye iron are deposits. The Tekovlevskoye ore strates now being existed is 10 km long, about 220 m wide. Its thickness varies from a few meters to 350 m and it has about 1,500 million tons of 61.4% rich iron-ore. There are 6 wet strata which will give 5,000 to 6,000 cu m of

water per hour when actual exploitation start. The authors say that the scheduled annual output is 15 million tons of ore. The mean exploitation coefficient will be 20.2 t/m2/year. The floors will sink

Card 1/3

APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-00513R000618610015-31

507/127-59-2-3/21

Preliminary Considerations Concerning the Methods of Opening, and Ways of Mining in the Yakovlavakove Deposit Area 10

> by about 6.9 m per year. The deposits will be exhausted in about 50 years. The authors defend the plans and advice of the Yuzhgiproruda Institute as opposed to the projects elaborated by the Institut gornogo dela AN SSSR (Institute of Mining attached to the Soviet Academy of Sciences). They especially argue against adapting the one-shaft-complex plan advocated by the Academy of Sciences. The proposed floor height is 70 to 80 m. The first 40% of the ore deposits are to be mined within 25 years, the next 27% within a further 14 years. A description and illustration of the actual preparatory work in the mines follows. Miner's trucks run by electric motors will each have 25 tons capacity. As far as the actual exploitation is concerned, the authors particularly recommend the self-collapsing floor system. Drainage operations will be carried out in 3 stages: deep-working pumps will first discard the pressure

Card 2/3

SOV/127-59-2-3/21


Preliminary Considerations Concerning the Methods of Opening, and Ways of Mining in the Kahovlevskope Deposit Area

of the subsoil waters; 2) a ring of drain shafts and galleries will be cut around the carbon limestone stratum; 3) then the ore layers will be drained. The floors placed at the bottom of the deposit must be equipped with a pumping system delivering 100 or 200 cu m of water per hour. There are 3

ASSOCIATION: Yuzhgiproruda, Khar'kov

Card 3/3

CIA-RDP86-00513R000618610015-3" APPROVED FOR RELEASE: 08/10/2001



INFANT'YEV, A.N., inch.

Questions of principle in opening thick, deep-lying deposits of rich iron ores in the Kursk Magnetic Anomaly. Izv.vys.ucheb.zav.;gor.zhur. 7 no.7:23-27 '64. (MIRA 17:10)

l. Yakovlevskiy rudnik Belgorodskoy oblasti. Hekomendovana kafedroy razrabotki rudnykh mestorozhdeniy Instituta gornogo dela.


IMENITOV, V.R., prof., doktor tekhn. nauk; CHIAYEV, T.I., gormyy inzh.; IMFANT'YEV, A.N.

THE CONTROL OF THE PROPERTY OF

Investigating the behavior of sand and clay depositions in the mining of iron ore deposits in the Kursk Magnetic Anomaly.

Gor. zhur. no.9:22-23 S 64. (MIRA 17:12)

1. Moskovskiy institut radioelektroniki i gornoy elektromekhaniki (for Imenitov, Chiayev). 2. Direktor Yakovlevskogo rudnika Kurskoy magnitnoy anomalii (for Infant'yev).



INFANT'YEV, VI. I. Cand Agr Sci — (diss) "Types of apple tree plantings in to Dzhunfar Ala-Tau , their natural restoration, growth, and productivity." Alma-Ata, 1957. 18 pp; Jaluftulle 19 cm. (Min of Agr USSR. Kazakh State Agr Inst). 100 copies. (KL, 22-57, 106).

-26-

INFANT YEV, V.I.

USSR / Forestry. Dendrology.

K-2

Abs Jour: Ref Zhur-Biol., No 6, 1958, 24877.

: Mushegyan, A. M.; Gribanov, L. N.; Infantiev, V.I.

Inst : Not given.

: On the Methods of Forest Valuation of the Haloxy-Title

lons of Kazakhstan.

Orig Pub: Lesn. kh-vo, 1957, No 8, 33-36.

Abstract: The exceptionally and increasingly difficult determination of the usual forest valuation indices of haloxylons is pointed out. It is proposed to divide the plantings into the following age groups: saplings, those ripening and those ripe. The criteria of the plantings of the black haloxylons ought to be established according to the proposed local table of the criteria, compiled on the basis of 120 test areas. A table to determine reserves

Card 1/2

26

CIA-RDP86-00513R000618610015-3" APPROVED FOR RELEASE: 08/10/2001

USSR / Forestry. Dendrology.

Abs Jour: Ref Zhur-Biol., No 6, 1958, 24877.

Abstract: of the haloxylon plantings according to the criteria, the average diameter at the surface of the ground and the degree of denseness of the plantings, is suggested.


Card 2/2

KARKLINS, J.; LIEPA, E.; INFANTYEVS, B.


a 1. de de la company de l La company de la company d

Latvijas Valsts universitates Zinatniskie raksti (Transactions of the Latvian State University); a review of Vols. 11-16. Vestis latv ak no.9:191-196 *59. (EEAI 9:10)

(Latvian periodicals)
(Academy of Sciences of the Latvian S.S.R.)



APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-00513R000618610015-3"



DEMIANSKI, M.; F INFEID, E.

Wote on the field method of obtaining the conservation laws and solving the two body problem in general relativity. Bul Ac Pol Mat 9 no.9:693-696 '61.

1. Institute of Theoretical Physics, University, Warsaw and Trinity College, Cambridge. Presented by L.Infeld.

DEMIANSKI, Marek; INFELD, Eryk


The field method of obtaining the conservat on laws and the Lagrangian. Acta physica Pol 21 no.5:469-479 My '62.

1. Univergity of Warsaw and Trinity College.

DEMIANSKI, M.; INFELD, E.

The radiative energy and the motion of particles. Bul Ac Pol mat 11 no.4:223-226 '63.

1. Institute of Physica, University, Warsaw, and Institute for Nuclear Research, Warsaw. Presented by L. Infeld.



INFIID, E.

Some exact solutions of the equations of magnetohydrodynamics for magnetic plane-symmetrical fields. Bul Ac Pol mat 12 no.4: 233-238 '64.


1. Institute of Nuclear Research, Warsaw. Presented by M. Danysz.

INFELD, Leopold The equations of motion in general relativity theory and the action principle. Acta physica Pel 16 no.3:177-210 *57. 1. Instytut Fizyki, Polska Akademia Nauk, Warszawa.


INFELD, Leopold


On studies of young scholars abroad. Nauka polska 10 no.3:91-93 My-Je 162.


Czlonek rzeczywisty Polskiej Akademii Nauk, Warszawa.





APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-00513R000618610015-3"













"APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3





Poland/Theoretical Physics - Quantum Electrodynamics

B-5

Abst Journal; Referat Zhur - Fizika, No 12, 1956, 33764

Author: Infeld, L., Plebanski, J.

Institution: University of Warsaw

Title : Electrodynamics Without Potentials

Original

Periodical: Acta Phys. Polon., 1953, 12, No 2, 123-134, English

Abstract: A general scheme was obtained for formulating a single theory of the electromagnetic field, characterized by an antisymmetric tensor $p_{\alpha,\beta}$ (interpreted as D and H) The simplest vector in this case will be $p^{\alpha\beta} = (4\pi/c)j^{\alpha}$. This equation is considered as a definition for the current. The Lagragian function H is considered in general as being dependent on the invariant $p = -1/4 p_{\alpha\beta} p^{\alpha\beta}_{j,\beta}$ and on the quantity $\rho = k(g_{\alpha\beta} p^{\alpha\beta}_{j,\beta} p^{\beta\mu}_{j,\beta})^{1/2}$, where k is some constant. The variational principle leads to

a field equation

Card 1/2

医克尔拉氏表现皮肤炎症或长足迹症的皮肤皮肤皮肤 医后槽 网络斯斯巴斯巴斯斯巴斯斯巴斯斯巴斯

Poland/Theoretical Physics - Quantum Electrodynamics

B-5

Abst Journal : Referat Zhur - Fizika, No 12, 1956, 33764

where $f_{\alpha\beta}=-2\partial H/\partial p^{\alpha\beta}$ is interpreted as the vectors E and B, while $A_{\alpha}=(c/4\pi)\partial H/\partial j^{\alpha}$ is considered to be the potential resulting from the theory. The energy-momentum tensor has the form


 $T_{\beta}^{\alpha} = 1/4[(H + 1/2 p^{\nu\mu} f_{\nu\mu})\delta_{\beta}^{\alpha} - p^{\alpha\nu} f_{\beta\nu}] + (1/c)[A_{\beta}j^{\alpha} - A_{\nu}j^{\nu}\delta_{\beta}^{\alpha}],$


which satisfies the equation $T_{\alpha,\beta}^{\alpha}=0$; all these quantities should be expressed in terms of $P_{\alpha,\beta}$ and their derivatives. It is shown that the proposed "Electrodynamics Without Potentials" is equivalent to the electrodynamics by Mie (Mie, G., Ann. Phys., 1912, 37, 511) in which the potentials are principal quantities. However, though the new Dirac electrodynamics can be formulated "without potentials" (Lagrangian $P + \rho/c$), it is outside the scope of the Mie electrodynamics.


Card 2/2

APPROVED FOR RELEASE: 08/10/2001

CIA-RDP86-00513R000618610015-3"








INFELD, L.

"Equations of Motion and Nonharmonic Coordinate Conditions," Byul, Polskoy akad. nauk, otd. 3,2, No 4, pp 161-164, 1954

The role of coordinate conditions in derivation of equations of motion of masses in a weak gravitational field is clarified. A transformation of coordinates, changing the field into a strong one, is always possible. The Newtonion equation of motion may be obtained from Einstein's equation as a first approximation, provided the gravitational field is weak and the motion quasistationary. (RZhFiz, No 6, 1955)

Sum. No. 681, 7 Oct 55



INFELD, L.

Einstein; reminiscent sketches, p. 349. (POSTEFY FIZYKI, Werszawa, Vol. 5, no. 3, 1954.)


SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. 5, Jan. 1955, Uncl.


INFELD, L.

Role of the theory of relativity in science, p. 355. [POSTEFY FIZYKI, Warszawa, Vol.5, no. 3, 1954.)

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. h, Jan. 1955, Uncl.

INFELD, Leopold. To Albert Einstein on his 75th birthday. Biul. VFER no. 10: 245-248 Ag-0'54. (NCRA 6:2) 1. Chlen Ispolnitel'nego kemiteta Vesmirnoy federatsii nauchnyth rabotnihov. (Einstein, Albert, 1879 - 1955)





INFELD, L.

From Copernicus to Einstein. p. 209. Vol. 1, no. 3, 1955. Warssawa

SERIA B: PRZYROD A NEOZYWIONA

SOURCE: East European Acession List (EEAL) Library of Congress Vol. 5, no. 8, August 1956

IMFELD, L.

Plebanski, J. Unitary transformations and spinor calculus. In English. p. 95. EULLETIN, Varsovie, Vol. 3, no. 2, 1955.

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, no. 10, Oct. 1955, Uncl.

INFEL D. L.

Category : USSR/Theoretical Physics - Quantum Field Theory

B-6

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 2951

Author : Infel'd, L.

: Institute of Theoretical Physics, Polish Academy of Science Inst

Title : Equations of Motion for Linear Field Theories

Orig Pub : Byul. Pol'sk. AN, Otd. 3, 1955, 3, No 4, 211-214

Abstract : It is noted that the equations of motion result from the field equations if the latter are nonlinear; in the case of linear equations, this does not take place. However, it becomes possible to derive the equations of motion from the field equations if the equations of the gravitational field are added to the system of linear equations. For example, in the case of the electromagnetic or meson field, it is necessary to start out with a system consisting of the

following equation Run- Igus R = - 8Th (Mas + Exs)

and the equations of the electromagnetic or meson field. Here Make and Edd are the tensor energy-momentum densities of the moving particles and R is the electromagnetic (or meson) field re-

Card : 1/2

APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-0 Category: USSR/Theoretical Physics - Quantum Field Theory CIA-RDP86-00513R000618610015-3"

. Abs Jour : Ref Zhur - Fizika, No 2, 1957 No 2951

The equations of motion are obtained for point masses in the following form:

(AT/AS) (MAN + EMN), A(3) X = 0

In the presence of only a single particle M assumes the form: M = $\{f(x), f(x)\}$ where m is the mass of the particle, $\{f(x)\}$ is the three-dimensional Dirac function; the dot denotes differentiation with respect to $\xi = t$. In the cartesian coordinate system, it follows from (1) that $m = m_0 \frac{dx}{ds}; \frac{dm_0}{ds} = -\frac{dx}{ds} \frac{d\xi_0}{ds} \int E_{,j} \int d\zeta_3 \times d\zeta_3$

so that in general the invariant mass \mathbf{m}_{Ω} is a function of the intrinsic time s.

Card : 2/2


Equations of motion.
Lecture delivered on 11th October, 1954 in Berlin during a celebration of the centenary of Riemann's work.


80: Progress in Physics, Poland, Vol. 6, #2, 1955, Unclassified.


INFELD, L. History of the theory of relativity. p. 96.


Vol. 6, no. 4, July 1955
FIZIKAI SZEMLE
SCIENCE
HUNGARY

So: East Europeon Accessions, Vol. 5, No. 9, Sept. 1956









APPROVED FOR RELEASE: 08/10/2001 CIA-RDP86-00513R000618610015-3"

INFELD, LEOPOLD

Moje wspomnienia o Einsteinie. Warszawa, Iskry, 1956. 148 p.

SOURCE:


Rast European Acession List (EEAL) Library of Congress Vol. 5, no. 8, August 1956


EINSTEIN, Alfred; LEFELD, Legold; SIMODOV, S.O.[translator]; LMSHKOVTSEV, F.A., redaktor; LIVSHITS, B.L., redaktor; TUMARKIMA, IW.A.tékhmichékikiy redaktor


[The evolution of physics; the growth of ideas from early concepts to relativity and quanta. Franzlated from the English Evolution is fished; resvited that of pervonsional hyper pointain do teorii otnositel nosti i kwant. Farevod a anglinkogo so vatup: stat'ei S.O. Suvorova. Ind. 2-oe. Moskva, Gos. ind-vo tekhniko-teoret. lit-try, 1956. 279 p. (MLRA 10:4)


(Physics-History) (Relativity (Physics))

(Quantum theory)





























INFELD

POLAND/Theoretical Physics - General

B-1

Abs Jour : Ref Zhur - Fizika, No 5, 1958, No 9892

Author : Infeld L., Plebanski, J.

Inst : Institute of Physics, Polish Academy of Sciences; The Univer-

sity, Warsaw, Poland.

: On a Further Modification of Dirac's -Functions Title

Orig Pub : Bull. Acad. polon. sci., 1957, Cl, 3, 5, No 1, 51-54

Abstract : Continuing their earlier work (Ref Zhur Fizika 1957, No 11, 27002), the authors introduce a three-dimensional (x) function of a new type, which satisfies the condition

[D(0) (()) x (x) [x] - = wp(P=1, x ..., K) where $J_{\omega}(0)$ is an arbitrary vicinity of the point $x = 0, \omega_0 = 1$, and w_p are pre-assigned numbers. An example of a \hat{j} (x)-function of this tpye is given.

Card : 1/1

INFELD, L.

APPROVED FOR RELEASE 108/100/10061ed Field-RDP86-00513R000618610015-3

Abs Jour : Ref Zhur - Fizika, No 3, 1958, No 5132

Institute of Physics, Polish Academy of Sciences : Infeld. Author : On the Lagrangian in Special Relativity Theory Inst

Title

Orig Pub : Bull. Akad. polon sci., 1957, Cl. 3,5, No 5, 491-495

Abstract : The relativistic equations of motion are derived from the variational principle. For this purpose the Lagrangian is chosen in the form $L^* = L(x_{,L}, x_{,L}) + (1/2)\gamma (x_{,L}, x_{,L})$ ds.

1), where γ is a certain scalar function, $x_{,L}$ dx.

By varying the action integral independently with respect to xy and y and then eliminating y it is possible to obtain the following Euler-Lagrange equations:

$$\left(\frac{\partial L}{\partial L}\right) - \left(\frac{\partial L}{\partial L}\right) - \left(\frac{\partial L}{\partial L} \times_{p}^{p} \times_{p}\right) + \left(L \times_{r}^{r}\right) = 0$$

: 1/2 Card

PHASE I BOOK EXPLOITATION

POL/4355

Polskie towarzystwo matematyczne

Prace Matematyczne, Seria I, II, 2 (Mathematical Transactions, Series I, vol. II.2)
Warszawa, Państwowe wyd-wo naukowe, 1958. 195 p. Errata slip inserted.
1,000 copies printed.

Editorial Board: WZadysław Orlicz (Chief Ed.), Stefan Drobot (Deputy Chief Ed.), Adam Bielecki, Stanisław Hartman, Jan Mikusiński, Roman Sikorski, Marceli Stark, Hanna Szmuszkowicz, Krzysztof Tatarkiewicz, and Wodzimierz Wrons.

FURPOSE: This book is intended for mathematicians

COVERAGE: The Contains 14 articles dealing with algebra, the theory of games, analysis, geometry, and two general mathematical topics. Summaries appear in Russian and English. No personalities are mentioned. References accompany individual articles.

Card 1/3

Mathematical Transactions (Cont.)	POL/4355
TABLE OF CONTENTS:	
	des for the Development
Infeld, L. (Warsaw). Importance of Modern Physics	201
Turski, S. (Warsaw). Mathematical Methods in Mc	dern Engineering 217
Steinhaus H. (Wrocław). McKinsey's Book on Gene	234
Sadowski, W. (Warsaw). Statistical Decision Fo	motions and the
Theory of Games	255
Straszewicz, S. (Warsaw). On the Basic Theorems	3 1n
Lobachevskiy's Trigonometry	269
Wilniewczyc, R. (Wroczaw). The Geometry of a Gr Transformations for Which A Parabola Rema	ins Invariant 287
Krasnodebski, R. (Wrockew). The Differential In	
Curve in a Symplectic Space	299
Mrówka, S. (Warsaw). On the Topology of the Spe	ices of
Closed Subsets	309
Mostowski, A. Włodzimierz (Warsaw). The Direct	Sums of Cyclic
Groups	319
Anna 0/2	
Card 2/3	

Mathematical Transactions (Cont.)		
Meder, J. (Szczecin). Application of the Mazur Theorem on	4355	
rzywicka, E. (Wood)	329	
olubian H on The Conditions at	:	
TO THE TAX THE TAX TO	337	
Delaber of Target Age	352	
Lesiekski, Z. (Poznań). On Some Inequalities ports From Scientific Sessions of the Polish Mathematical Society AHABIE: Library of Government	355 361	
AHABIE: Library of Congress	368	
rd 3/3		
AC/dva 10/19/	/in 60	
		Ř

INFELD, L.

My reminiscences of Wladyslaw Natanson.

p. 3. (KOSMOS. SERIA B: PRZYWODA NIEOZYWIONA.) (Warszawa, Poland) Vol. 4, no. 1, 1958

SO: Monthly Index of East European Accession (EEAI) LC Vol. 7, No. 5, 1958

INFELD, L.

The genealogy of Sputnik.

p. 9. (KOSMOS. SERIA B: PRZYWODA NIEOZYWIONA.) (Warszawa, Poland) Vol. 4, no. 1, 1958

SO: Monthly Index of East European Accession (EEAI) LC Vol. 7, No. 5, 1958

INFELD, L.

"Planck's hundredth anniversary"

p. 205 (Kosmos, Seria B; Przyroda . Nieczywiona, Journal on natural sciences with the exception of biology issued by the Copernicus Society of Polish Naturalists, Vol. 4, no. 3, 1958, Warsaw, Poland)

Monthly Index of East European Accessions (EEAI) LC, Vol. 8, No. 1, Jan. 59.

POLAND/Nuclear Physics - General Problems.

: Ref Zhur Fizika, No 2, 1960, 2726

Author

Abs Jour

Infeld, Leopold

Inst

Title

: Impressions of the Second International Conference on

Atomic Energy in Geneva

Orig Pub

: Kosmos (Polska), 1958, B4, No 4, 273-275

Abstract

No abstract.

Card 1/1

CIA-RDP86-00513R000618610015-3" APPROVED FOR RELEASE: 08/10/2001

POLAND/Nuclear Physics - Physical Base of Nuclear and

Thermonuclear Technology.

Abs Jour

: Ref Zhur Fizika, No 1, 1960, 614

Author

Infeld, Leopold

Inst Title : Impressions of the Second Conference on Atomic

Energy in Geneva

Orig Pub

: Nukleonika, 1959, 4, No 1, 1-4

Abstract

: No abstract.

_ INFELD, L.

A new form of the geodesic line equation. Bul Ac Pol mat 8 no.8: 559-561 *60.

1. Institute of Physics, University, Warsaw and Institute of Physics, Polish Academy of Sciences.

(Geodesy) (Equations)

INFELD, L.

SURTIAME (in caps); Given Names

Country:

Poland

Academic Degrees: Not stated

Institute of Physics (Instytut Fizyki), Polish

Affiliation:

Academy of Sciences (Polska Akademja Nauk)

Source:

Warsaw, Bulletin de l'Académie Polonaise des Sciences,

Série des Sciences Mathématiques, Astronomiques et

Physiques, Vol 9, No 2, Feb 61, pp 93-97.


Data:

The EIH and the k-Approximation Methods."


CIA-RDP86-00513R000618610015-3" APPROVED FOR RELEASE: 08/10/2001


On the most Cartesian-like coordinate system. Bul Ac Pol Mat 9 no.4: 299-302 '61.


1. Institute of Theoretical Physics, Warsaw University.



INFELD, Leopold; NACY, Tibor [translator]
Gravitation, 1962. Fiz szemle 12 no.11:354 N '62.










INFELITSYN, A., (Engr-Lt Col)

Coauthor with Engr-Lt Col. I. CHEPELEVSKIY* of article, " Tent for Repairing Equipment," concerning the construction of a tent to be used in the field when repairing equipment. (Tankist, Moscow, No 4, Apr. 1954)

SO: SUM No 239, 13 Oct. 1954









INGARDEN, R.S.

Equations of motion and field equations in five-dimensional unified relativity theory. Dokl. Akad. Nauk SSSR 88, No.5, 773-6 53. (MLRA 6:2) (PA 56 no.671:7416 53)

States that in the theory of relativity there exist two different methods for deriving the eqs of motion from the field eqs; the method of Einstein and Infeld on the one hand and the method of Fok on the other. Attempts to show that these two views can agree to a certain extent in a 5-dimensional "unified" theory of relativity, in which a new point of view is given to the problem. Presented by Acad V. A. Fok 20 Dec 52. Indebted to V.A. Fok for his helpful remarks made at the configurace of Polish physicists at Spala.





LN9ARDEN R.S.
POLAND/Optics - Optical Technology

K-4

Abs Jour : Ref Zhur - Fizika, No 4, 1958, No 9157

Author : Ingarden, R.S., Okhman, G.

: Mathematics Institute, Academy of Sciences, Warsaw, Inst

Title : Optimum Optical Systems

Orig Pub : Syul. Pol'skoy AN, otd. 3, 1954, 2, No 6, 275-280

Abstract : Determination of a criterion that characterizes a system with

the best image quality. Systems are considered with axial symmetry, consisting of homogeneous and isotropic media. For the sake of simplicity, non-self-illuminating objects are taken, and the investigation is carried out in the meridional plane. The action of an optical system is represented, using Mandel'shtam's example, as an integral equation that transforms the amplitude in the plane of the object into an amplitude in the plane of the image, the kernel of which depends only on the optical system. It is shown that an optical system having no aberration is not ideal from the point of view of the wave theory of light. Only a system satisfying definite conditions

will reproduce the object with absolute similarity.

Card : 1/2

INCARDEN, R.

Bulletin - Vol. 2, no. 7, 1954.

Embedding Finsler spaces in a Minkowski space. p. 305.

SO: Monthly list of East European Accessions, (EEAL), LC, Vol. 4, No. 9, Sept. 1955
Uncl.